



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>A23L 2/62</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 95/34223</b> <b>(43) International Publication Date:</b> 21 December 1995 (21.12.95)
<b>(21) International Application Number:</b> PCT/DK95/00224 <b>(22) International Filing Date:</b> 7 June 1995 (07.06.95) <b>(30) Priority Data:</b> 683/94 15 June 1994 (15.06.94) DK <b>(71) Applicant (for all designated States except US):</b> NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsværd (DK). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> HELDT-HANSEN, Hans, Peter [DK/DK]; Novo Nordisk A/S, Novo Allé, DK-2880 Bagsværd (DK). HYTTEL, Susanne [DK/DK]; Dalgas Have 2, 3. t.v., DK-2000 Frederiksberg (DK). <b>(74) Common Representative:</b> NOVO NORDISK A/S; Corporate Patents, Novo Allé, DK-2880 Bagsværd (DK).		<b>(81) Designated States:</b> AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> EXTRACTS/CLOUD STABILITY  <b>(57) Abstract</b>  The invention relates to a method of producing cloud stable extracts such as juices from plant material by using one or more enzymes that attack the hairy regions of pectin.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgystan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

**EXTRACTS/CLOUD STABILITY****FIELD OF INVENTION**

This invention relates to a method of producing cloud stable extracts from plant material useful for producing juice  
5 and nectar.

**BACKGROUND OF THE INVENTION**

The trends in modern plant extract technology are moving towards the use of a greater variety of raw materials, a more complete utilisation of these raw materials, a speeding  
10 up of processes and a presentation of a greater variety of bases and finished products. Some of these developments are made possible through improvements of existing processes and process equipments, as well as through introduction of new processes and equipments. The use of enzyme preparations as  
15 processing aids play a key role in these developments.

During fruit juice manufacture enzyme preparations are often used in the steps of extraction and liquefaction of fruit and fruit juice clarification. The commercial enzyme preparations contain a mixture of enzymes which degrade the  
20 pectin polymers (including pectin lyases, polygalacturonases, pectin esterases, galactanases, arabinases), as well as other enzymes like cellulases and xylanases.

New developments are going into the direction of cloudy juices and extracts from plant material. The mechanism  
25 why some of these extracts stay cloudy while others spontaneously clarify is not well understood. Therefore emphasis is put into research to learn more about the background of cloudiness and cloud stability.

Pectins (pectin-polymers) occur in nature as  
30 constituents of higher plant cell walls. They are found in primary cell wall and middle lamella where they are embedded in cellulose fibrils. The composition of pectin is variable among plant species and moreover dependent on the age and the

**SUBSTITUTE SHEET**

maturity of the fruit. Among the richest sources of pectins are lemons and oranges, which can contain up to 30% of this polysaccharide.

Most pectin-polymers are composed of smooth regions, i.e. linear homogalacturonan, and hairy (ramified) regions.

The linear homogalacturonan is composed of chains of 1,4-linked  $\alpha$ -D-galacturonic acid, the polygalacturonic acid is methoxylated to a varying extent, and may further be partially acetylated. The linear homogalacturonan can be degraded and depolymerized by different enzymes: Pectin lyase which cleaves the galacturonosyl bonds of highly methoxylated pectins by  $\beta$ -elimination. Pectate lyase cleaves galacturonosyl bonds in the non-methoxylated parts of pectin by  $\beta$ -elimination, and polygalacturonase hydrolyses the glycosidic linkages in the non-methoxylated part of homogalacturonan. The action of pectate lyase and polygalacturonase is facilitated by pectin methyl-esterase which catalyses the removal of methanol from homogalacturonan, resulting in the formation of pectic acid (polygalacturonic acid). Enzymes or enzyme combinations which have the ability to depolymerize homogalacturonan are designated homogalacturonan depolymerizing enzymes in the following disclosure.

The hairy regions consist of a rhamnogalacturonan backbone with side-branches of varying length. The pectin hairy regions might be heterogeneous, with regions with extensive branching, regions with less extensive branching, and regions where the backbone is rich in galacturonic acid with an extensive branching with  $\beta$ -linked xylose (xylogalacturonan).

The composition of the very complex structure of the hairy regions vary according to the source of the plant cell wall, cf. Schols et al. in Carbohydrate Research 206, 1990, pp. 117-129; O'Neill et al. in "Methods in Plant Biochemistry", Vol. 2, Carbohydrates, P. M. Dey (Ed.), 1990, Academic Press, London, pp. 415-441; Voragen and Schols in "Structural Studies of Plant Cell-Wall Polysaccharides Using Enzymes", Special Publication No. 134, The Royal Society of Chemistry 1993 and Carpita and Gibeaut in The Plant Journal 3(1), 1993, pp. 1-30.

Rhamnogalacturonans are polysaccharides with more or less regularly alternating rhamnose and galacturonic acid residues in the backbone. The rhamnogalacturonan backbone in the hairy regions has acetyl groups on the galacturonic acid residues (cf. H.A. Schols in Carbohydrate Research 206, 1990, pp.117-129).

The degradation of the backbone of the hairy regions is performed by enzymes designated rhamnogalacturonases (RGases). RGases are believed to hydrolyse the bond between rhamnose and galacturonic acid. In order to facilitate the activity of RGases it may be desirable to reduce the degree of acetylation of the backbone, e.g., by use of the enzyme rhamnogalacturonan acetyl esterase (cf. Searle-van Leeuwen et al. in Appl. Microbiol. Biotech. 38, 1992, p. 347-349). Furthermore, a reduced degree of branching of parts of the hairy regions may facilitate the activity of rhamnogalacturonases. The reduced degree of branching may be obtained by enzymes which attack the side-branches.

The isolation and purification of a RGase from Aspergillus aculeatus are described by Schols et al. in Carbohydrate Research 206, 1990, p. 105-115. Another type of RGase from A. aculeatus is described in WO 92/19728.

Enzymes which attack the backbone of hairy regions are defined in the following disclosure as any enzyme or combination of enzymes which has the capability of attacking (by hydrolysis,  $\beta$ -eliminations, or otherwise) the backbone of hairy regions, e.g., rhamnogalacturonase containing enzymes preparations.

The side-branches include monosaccharides like xylose, galactose and arabinose, and oligo and polysaccharides like arabinan, galactan and arabinogalactan.

Galactan contains  $\beta$ -1,4 linked galactose in the backbone. Galactans are degraded by  $\beta$ -1,4-galactanases (EC 3.2.1.89) (in short galactanases). Reference can be made to R.F.H. Dekker and G.N. Richards, "Hemicellulases, their Occurrence, Purification, Properties and Mode of Action" in R.S. Tipson and D. Horton, Advances in Carbohydrate Chemistry and

Biochemistry, Academic Press 32, 277-352 (1976); R.F.H. Dekker, "The Hemicellulase Group of Enzymes", in J.M.V. Blanchard and J.R. Mitchell, Polysaccharides in Food, Butterworths, 93-108 (1979), and A.G.J. Voragen, F. Geerst and W. Pilnik "Hemicellulases in Enzymatic Fruit Processing", in P. Depuy, Use of Enzymes in Food Technology, Technique et Documentation Lavoisier, 497-502 (1982). One example of a galactanase is the galactanase described in WO 92/13945.

Further, galactan and galactose sidebranches are degraded by the exo-acting enzyme  $\beta$ -galactosidase.

Galactans might have arabinose sidebranches (arabinogalactan), these sidebranches are hydrolysed by  $\alpha$ -arabinosidase. The partial or full removal of the arabinose sidebranches might facilitate the activity of galactanases.

Arabinan is composed of a backbone of  $\alpha$ -L-arabinose subunits linked  $\alpha$ -(1 $\rightarrow$ 5) to each other and side chains linked  $\alpha$ -(1 $\rightarrow$ 3) or  $\alpha$ -(1 $\rightarrow$ 2) to the backbone. Enzymes which are capable of degrading arabinan backbone are designated arabinanases. The sidebranches of arabinan can be hydrolysed by  $\alpha$ -arabinosidases (Rombouts et al., Carbohydrate Polymers 9, 1988, p. 25), which can also hydrolyse linear arabinan from the non-reducing end.

Xylose side-branches can be removed by  $\beta$ -xylosidase.

Enzymes which attack the hairs of the hairy regions of pectin will in the following disclosure include all enzymes which have the ability to degrade fully or partially the hairs of the hairy regions of pectin, including galactanase,  $\beta$ -galactosidase,  $\beta$ -xylosidase, arabinanase and  $\alpha$ -arabinosidase, or any combination hereof.

The term hairy region degrading enzymes are meant to include both enzymes which attack the hair or the backbone of the hairy regions.

The enzyme preparations used for production of plant extract like juice contain both homogalacturonan depolymerization enzymes as well as other enzymes like enzymes which attack the hairy regions of pectin and/or cellulose. Such enzyme combinations often lead to extracts with a low cloudiness and/or a low cloud stability. Homogalacturonan depolymer-

rization activity can be measured as PSU activity (Analytical method obtainable from Novo Nordisk A/S as AF-269).

#### SUMMARY OF THE INVENTION

It is an object of the present invention to provide  
5 a method of producing cloudy extracts with improved cloud stability from plant material. Surprisingly, it has been found that the use of taylormade monocomponent enzyme preparations is a key function to reach this goal.

Accordingly, the invention provides a method of  
10 producing cloud stable extracts from plant material comprising use of one or more enzymes that attack hairy regions in the plant material.

#### DETAILED DISCLOSURE OF THE INVENTION

##### Extracts

15 In the context of this invention an extract from a plant material is any substance which can be derived from plant material by extraction, processing or by other separation techniques. The extract may be juice, nectar, base, or concentrates made hereof. The plant material may be vegetables,  
20 e.g., carrots, celery, onions, or fruits, e.g., pome or seed fruits (apples, pears, etc.), grapes, tomatoes, citrus (orange, lemon, lime, mandarin), prunes, cherries, black currants, red currants, raspberries, strawberries, cranberries, pineapple, all kinds of tropical fruits. According to this invention  
25 especially juice from apples is preferred.

##### Enzymes

In order to produce cloudy extracts and/or extracts with an improved cloud stability, different enzyme(s) or a combination of different enzymes may be used. It is preferred  
30 to add an enzyme which attacks the hairs of the hairy regions of pectin such as a galactanase, an arabinanase, a  $\beta$ -galac-

tosidase, a  $\beta$ -xylosidase and/or an  $\alpha$ -arabinosidase, e.g., a galactanase obtained as described in WO 92/13945.

Additionally, an enzyme which attacks the backbone of the hairy regions, e.g., a rhamnogalacturonase, preferably a rhamnogalacturonase II (obtained as described in WO 92/19728) may be added, added alone or together with one or more enzymes that attack the hairs of the hairy regions. To further increase the cloud stability of the extract a rhamnogalacturonan acetyl esterase (which may be obtained as described in WO 93/20190) may be added. Additionally enzymes which attack other parts of the plant material can be added, like xylanases and endoglucanases.

Preferably, the enzyme preparations should be free from homogalacturonan depolymerizing ability, as such enzymes are seen to reduce the cloudiness and cloud stability which can be obtained by the hairy region degrading enzymes.

#### Cloudiness

The cloudiness of a plant extract can be measured as the turbidity, e.g., by a Nephla Turbidity Photometer conforming to DIN 38404 and ISO 7027 using a formazin DIN standard.

#### Cloud Stability

According to this invention the stability  $\Delta T_z(\%)$  of the cloud particles is determined by the turbidity of the supernatant of the extract after centrifugation at 4160 X g, for 15 minutes relative to the turbidity of the un-centrifuged extract. The forced precipitation of the cloud particles in a given extract is an indication for the stability of the continuous phase to keep the particles in dispersion.

More specifically the method is based upon a centrifugation of a 60 ml extract sample in a glass centrifuge tube and centrifuged for 15 min at 4160 X g. The turbidity before ( $T_0$ ) and after centrifugation ( $T_z$ ) is measured by a Nephla Turbidity Photometer conforming to DIN 38404 and ISO 7027 using a formazin DIN standard. The cloud stability  $\Delta T_z(\%)$  is then calculated as:

SUBSTITUTE SHEET



$$\Delta T_z(\%) = [(T_z)/(T_0)] \times 100.$$

The cloudiness and cloud stability of a plant extract like apple juice vary significantly according to the sort of the apple and the maturity of the apples. The enzyme(s) in accordance with this invention will increase the cloudiness and cloud stability relative to what can be obtained without the use of enzymes.

The enzymes in accordance with the invention may be used in a process where the plant material is milled, chopped, or otherwise fragmented. The fragmented plant material is then treated with the enzymes for 1 min to 48 hours, preferably 10 min to 8 hours, most preferably 30 min to 4 hours, at 10 - 60°C, preferably 15 - 50°C, most preferably 15 - 40°C. The extract can then be obtained by pressing, decanting, centrifugation or by other separation techniques. The extract may be further processed e.g by enzymes, by filtration, by concentration or otherwise.

The invention is further illustrated in the following example, which is not intended to be in any way limiting to the scope of the invention as claimed.

SUBSTITUTE SHEET

**EXAMPLE 1**Cloud Stable Apple Juice

Cloud stability was tested in three apple sorts (Jona Gold, Mutzu and Belle de Boskop) using different enzyme combinations. In each apple test the following procedure was used:

The apples were stored for at least 12 hours at room temperature to make sure that the apples had a temperature suitable for incubation with the enzymes in question.

10        1.1 kg of apples were cut in four parts, and milled in a vegetable cutting machine (Brøttner type R 301 equipped with a 1.5 mm grater). Ascorbic acid was added to the mash to obtain a concentration of 0.1% to the mash to prevent oxidation. Then the different enzyme combinations were added (the  
15 combinations described below) in an amount of 25 mg of pure enzyme protein of each enzyme per kg of mass. The different enzymes were added simultaneously to the mash (the amount of pure enzyme protein was estimated by SDS-PAGE; the band on the SDS-PAGE gel, which correspond to the enzyme in question, can  
20 be identified by purifying the enzymes by methods known in the art).

The mash was enzyme treated for 2 hours at room temperature, whereafter the mash was processed by a HAFICO press machine at 300 kg/m<sup>2</sup> for 5 min. Then the freshly pressed  
25 apple juice was pasteurized (85 - 90°C, 15 min.), cooled to room temperature (20 - 25°C), and the cloudiness and cloud stability were measured as described above.

The following enzymes were used in the experiments:

Galactanase I obtained as described in WO 92/13945.

30 Rhamnogalacturonase II obtained as described in WO 92/19728.

Rhamnogalacturonan acetyl esterase as described in WO 93/20190.

Endoglucanase III. Endoglucanase III is equivalent to the endoglucanase (FI-CMCase) described by T. Ooi et al in Nucleic Acids Res. 18, 1990, p. 5884, and in Biotech. Biochem. 57,

35 1993, p. 1960-1961, cloned and expressed as described in the

SUBSTITUTE SHEET

Danish Application DK 0245/93, using the cellulase detection method described in WO 93/11249.

Polygalacturonase as described in the Danish application DK 1545/92.

5 Pectin methylesterase as described in the Danish application DK 487/93.

4 separate experiments were carried out. All the enzymes compared in each experiments were tested on each of three experimental days.

10 Experiment 1:

Apple sort: Jona Gold:

Tested enzymes:

1) Galactanase I (Gal I)

2) Gal I + Rhamnogalacturonase II (Rham II) + Rhamnogalac-  
15 turonan acetyl esterase (Rham Ac)

3) Gal I + Rham II + Rham Ac + Endoglucanase III (End III)

The result are presented in Table 1.

Experiment 2:

Apple sort: Red Belle de Boskop

20 1) Gal I

2) Gal I + Rham II + Rham Ac

3) Gal I + Rham II + Rham Ac + End III

The results are presented in Table 2.

The same type of enzymatic effects are seen on the Jona Gold  
25 and Red Belle de Boskop (cf. Table 1 + 2). It is seen that the hairy region debranching enzyme galactanase increases the cloudiness as well as the cloud stability. These effects are markedly improved by further addition of the enzyme combination rhamnogalacturonase and rhamnogalacturonan acetyl esterase,  
30 which attack the backbone of the hairy regions. Addition of endoglucanase will even further improve the cloud stability.

SUBSTITUTE SHEET

Experiment 3:

Apple sort: Mutzu

- 1) Gal I
- 2) Gal I + Rham II + Rham Ac
- 5 3) Gal I + Rham II + Rham Ac + End III

The results are presented in Table 3.

It is seen from Table 3 that galactanase alone is not able to improve the cloudiness of the apple sort Mutzu. The combination of the debranching enzyme galactanase and the backbone degra-  
10 ding enzyme combination (rhamnogalacturonase and rhamnogalacturonan acetyl esterase) improves cloudiness and cloud stability. These quality parameters are further improved by endoglucanase.

Experiment 4:

15 Apple sort: Red Belle de Boskop

- 1) Gal I + Rham II
- 2) Gal I + Rham II + Rham Ac + Pectin methyl esterase (Pme) + Polygalacturonase (Polygal)

The results are presented in Table 4.

20 It is observed from Table 4 that the combination of galactanase and rhamnogalacturonase (without rhamnogalacturonan acetyl esterase) is able to give a strong improvement of cloudiness and cloud stability. Further it is demonstrated that the addition of a homogalacturonan depolymerising enzyme preparation  
25 tion of polygalacturonase and pectin methyl esterase, will destroy the cloud stability, even though the enzyme preparation also contains galactanase, rhamnogalacturonase and rhamnogalacturonan acetyl esterase, which otherwise will give a cloud stable juice.

Table 1  
Apple Variety: Jona Gold

Enzyme Combination	Experimental No. / Average	Turbidity T <sub>90</sub> before centrifugation	Increase in turbidity relative to untreated control (%), based on the average values	Cloud stability (( $\Delta T_z$ )(%))
Untreated	1	1056		33
	2	540		33
	3	897		35
	Average	831	100	34
Gal I	1	1144		42
	2	743		29
	3	1047		38
	Average	978	118	36
Gal I + Rham II + Rham Ac	1	1349		51
	2	1340		43
	3	1195		60
	Average	1295	159	51
Gal I + Rham II + Rham Ac + End III	1	1270		57
	2	1325		53
	3	1178		66
	Average	1257	151	59

SUBSTITUTE SHEET

Table 2  
Apple Variety: Red Belle de Boskoop

Enzyme Combination	Experimental No. / Average	Turbidity T <sub>90</sub> before centrifugation	Increase in turbidity relative to untreated control (%), based on average values	Cloud stability (( $\Delta T_z$ (%))
Untreated	1	1109	100	57
	2	932		58
	3	1140		53
	Average	1061		56
Gal I	1	1193	114	94
	2	1232		60
	Average	1212		77
Gal I + Rham II + Rham Ac	1	1423	125	86
	2	1222		95
	3	1354		76
	Average	1333		86
Gal I + Rham II + Rham Ac + End III	1	1273	117	81
	2	1143		93
	3	1323		86
	Average	1246		90

SUBSTITUTE SHEET

Table 3  
Apple Variety: Mutzu

Enzyme Combination	Experimental No. / Average	Turbidity $T_0$ , before centrifugation	Increase in turbidity relative to untreated control (%), based on average values	Cloud stability ( $\Delta T_2$ (%))
Untreated	1	1197	100	37
	2	1072		35
	Average	1134		36
Gal I	1	916	94	43
	2	1111		26
	3	1156		24
	Average	1061		31
Gal I + Rham II + Rham Ac	1	1235	112	48
	2	1250		54
	3	1337		35
	Average	1273		46
Gal I + Rham II + Rham Ac + End III	1	1156	113	47
	2	1268		68
	3	1420		55
	Average	1281		57

Table 4  
Apple Variety: Red Belle de Boskoop

Enzyme Combination	Experimental No. / Average	Turbidity T <sub>90</sub> before centrifugation	Increase in turbidity relative to untreated control (%), based on average values	cloud stability (( $\Delta T_2$ (%))
Untreated	1	558	100	4
	2	1375		61
	3	479		1
	Average	804		22
Gal I + Rham II	1	1315	172	87
	3	1463		54
	Average	1389		70
Gal I + Rham II + Rham Ac + Pme + Polygal	1	350	157	1
	2	691		2
	3	1307		0
	Average	1265		1

SUBSTITUTE SHEET



**CLAIMS**

1. A method of producing cloud stable extracts from plant material comprising use of one or more enzymes that attack the hairy regions of pectin.

5           2. A method according to claim 1, wherein the extract is juice, nectar, base or concentrate.

3. A method according to claim 1, wherein the plant material is obtainable from vegetables or fruits.

4. A method according to claim 3, wherein the fruits  
10 are selected from the group consisting of apples, pears, citrus, (orange, lemon, lime, mandarin), tomatoes, grapes, black currants, red currants, raspberries, strawberries, cranberries, prunes, cherries, pineapple and all kinds of tropical fruits.

15           5. A method according to claim 4, wherein the fruits are apples.

6. A method according to claim 3, wherein the vegetables are selected from the group consisting of carrots, celery and onions.

20

7. A method according to claims 1-6, wherein one or more enzymes attack the hairs of the hairy regions of pectin, preferably selected from the group consisting of galactanases,  $\alpha$ -arabinosidases,  $\beta$ -xylosidases,  $\beta$ -galactosidases and arabina-  
25 nases, most preferably selected from galactanases.

8. A method according to claims 1-6, wherein one or more enzymes attack the backbone of the hairy regions of pectin, preferably a rhamnogalacturonase.

9. A method according to claim 8, additionally comprising use of a rhamnogalacturonan acetyl esterase.

10. A method according to claims 1-7, additionally comprising use of one or more enzymes that attack the backbone  
5 of the hairy regions of pectin, preferably a rhamnogalacturonase.

11. A method according to claim 10, additionally comprising use of a rhamnogalacturonan acetyl esterase.

12. A method according to claims 1-11, additionally  
10 comprising use of an endoglucanase.

13. A method according to claims 1-12, wherein the enzymes are without significant homogalacturonan depolymerising activity.

14. A method according to claim 13, wherein the  
15 enzymes are without significant activity of pectin lyase, polygalacturonase or pectate lyase.

15. An enzyme preparation comprising at least one enzyme that attacks the hairs of the hairy regions of pectin and at least one enzyme that attacks the backbone of the hairy  
20 regions of pectin, and without significant homogalacturonan depolymerising activity.

18. An enzyme preparation according to claim 17 comprising galactanase and rhamnogalacturonase activity and without significant activity of pectin lyase, polygalacturonase  
25 or pectate lyase.

19. An enzyme preparation according to claim 18, additionally comprising rhamnogalacturonan acetyl esterase and/or endoglucanase.

<b>A. CLASSIFICATION OF SUBJECT MATTER</b>		
IPC6: A23L 2/62 According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b>		
Minimum documentation searched (classification system followed by classification symbols)		
IPC6: A23L		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
SE,DK,FI,NO classes as above		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
DIALINDEX (FOODSCI), EPODOC		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Journal of Food Science, Volume 43, 1978, P. STRUEBI et al., "Use of a macerating pectic enzyme in apple nectar processing" page 260 - page 263 --	1-5
X	Flüssiges Obst, Volume 58, No 5, 1991, A. ASKAR et al., "Polysaccharide und Trubstabilität tropischer Nektare" page 244 - page 251 --	1-4
X	Dialog Information Service, File 51, Food Sci. & Tech. Abs, Dialog accession no. 00343482, FSTA accession no. 87-09-h0172, M. Anastasakis et al. "Enzymatic hydrolysis of carrot for extraction of a cloud-stable juice", Food Hydrocolloids (1987), 1 (3) 247-261 --	1,3,6
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex.		
<ul style="list-style-type: none"><li>* Special categories of cited documents:</li><li>"A" document defining the general state of the art which is not considered to be of particular relevance</li><li>"B" earlier document but published on or after the international filing date</li><li>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</li><li>"O" document referring to an oral disclosure, use, exhibition or other means</li><li>"P" document published prior to the international filing date but later than the priority date claimed</li><li>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</li><li>"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</li><li>"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</li><li>"&amp;" document member of the same patent family</li></ul>		
Date of the actual completion of the international search		Date of mailing of the international search report
25 Sept 1995		28-09-1995
Name and mailing address of the ISA/ Swedish Patent Office Box 5055, S-102 42 STOCKHOLM Facsimile No. +46 8 666 02 86		Authorized officer  INGA-KARIN PETERSSON Telephone No. +46 8 782 25 00

2  
INTERNATIONAL SEARCH REPORT

International application No.  
PCT/DK 95/00224

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4388330 A (H.J. WOBEN ET AL.), 14 June 1983 (14.06.83), claims 1,8  -----	1-4

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

28/08/95

International application No.

PCT/DK 95/00224

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A- 4388330	14/06/83	DE-A,C,C 3045817	19/06/81
		JP-C- 1470758	14/12/88
		JP-A- 56088784	18/07/81
		JP-B- 63019148	21/04/88
		NL-A- 7908722	01/07/81
<hr/>			

Form PCT/ISA/210 (patent family annex) (July 1992)

**THIS PAGE BLANK (USPTO)**